

REVIEW ARTICLE

The Clinical Implications of Blood Adiponectin in Cardiometabolic Disorders

Lin-Chau Chang,¹ Kuo-Chin Huang,³ Yen-Wen Wu,^{4,5} Hsien-Li Kao,^{2,4} Chi-Ling Chen,¹
Ling-Ping Lai,^{2,4} Juey-Jen Hwang,^{2,4} Wei-Shiung Yang^{1,2,4*}

Adipose tissue is now accepted by the scientific and medical community to be a genuine endocrine organ, in addition to its classical role as an energy store. Adiponectin is one of the many adipocytokines that are secreted almost exclusively by adipose tissue. Alteration in blood adiponectin concentrations has been linked to many human diseases in numerous cross-sectional and prospective studies. In this review, we describe briefly the biological effects of adiponectin as revealed by basic scientific investigations. We also summarize the principles of blood adiponectin assays. Overall, lower blood adiponectin concentration is found in subjects with obesity, type 2 diabetes mellitus, dyslipidemia, and hypertension. These medical conditions are components of the metabolic syndrome and major risk factors for accelerated atherosclerosis. Plasma adiponectin levels are also expected to be lower in subjects with cardiovascular diseases, such as coronary artery disease, ischemic stroke and peripheral artery disease. Congestive heart failure (CHF) and cardiac arrhythmia are common end points in cardiovascular diseases. Surprisingly, higher blood adiponectin levels are frequently reported to predict mortality associated with CHF. Few human data regarding adiponectin and cardiac arrhythmia are available. Higher blood adiponectin level has been documented only in atrial fibrillation. We also summarize data on the role of the high molecular weight (HMW) isoforms of adiponectin and the effects of clinical treatment on the levels of total or HMW adiponectin. Whether adiponectin is a risk marker or a risk factor for the diseases reviewed in this article, and in many other human diseases, and their detailed pathogenic links awaits further investigation. [*J Formos Med Assoc* 2009;108(5):353–366]

Key Words: adiponectin, arrhythmia, atrial fibrillation, cerebrovascular disease, coronary artery disease, diabetes, dyslipidemia, heart failure, hypertension, metabolic syndrome, obesity, peripheral vascular diseases, stroke

Fat Hormone

Obesity has become a global epidemic in children and adults, and is considered to be the root cause of many metabolic diseases that we face today.¹ Through advancing our understanding of the pathogenesis of obesity, we no longer view adipose tissue as a simple fat reservoir for energy storage.

Instead, adipose tissue is now considered an endocrine organ, actively regulating energy balance and many other physiological functions.² The crucial evidence leading to the formation of this concept was the discovery of leptin.

More than 40 years ago, monogenic obesity and diabetes mouse models, obese (*ob*) and diabetes (*db*), were identified.^{3,4} These two models were

©2009 Elsevier & Formosan Medical Association



ELSEVIER

¹Graduate Institute of Clinical Medicine and ²Department of Medicine, College of Medicine, National Taiwan University; Departments of ³Family Medicine, ⁴Internal Medicine, and ⁵Nuclear Medicine, National Taiwan University Hospital, Taipei, Taiwan.

Received: February 25, 2009

Revised: February 28, 2009

Accepted: March 2, 2009

***Correspondence to:** Dr Wei-Shiung Yang, Department of Medicine, College of Medicine, National Taiwan University, 1 Chang-Teh Street, Taipei 100, Taiwan.
E-mail: wsyang@ntu.edu.tw

demonstrated to harbor two spontaneous recessive mutant alleles. Although they share similar syndromes of hyperphagic obesity with prominent diabetes and endocrine dysfunction, they are in fact caused by two different genes.⁵ The parabiosis experiments conducted by Coleman et al provided insight into the underlying mechanisms.^{6,7} Connecting the blood circulation of *ob/ob* mice with that of wild-type or *db/db* mice led to significant weight loss in the *ob/ob* animals, while connecting the circulation of *db/db* mice with that of *ob/ob* or wild-type mice exerted no effect on the *db/db* animals. These findings indicate that a circulating factor that causes weight loss is missing in *ob/ob* mice, while possibly the receptor for the factor is lacking in *db/db* mice. This circulating factor, a 16-kDa peptide hormone, was later discovered by positional cloning of the *ob* gene in mice in 1994.⁸ Since the factor promoted weight loss in *ob/ob* mice, it was termed leptin, which was derived from "leptos", the Greek for thin.⁹ Soon thereafter, the leptin receptor was also cloned in 1995.¹⁰ The leptin receptor is exactly what is missing in *db/db* mice. Leptin therefore became the first example of the fat hormones, or the so-called adipocytokines or adipokines, which are produced mainly by the adipose tissues and have significant systemic effects. A comprehensive review of leptin is beyond the scope of the present article. Readers interested in this are advised to read some recent reviews.^{11,12}

Among the adipocytokine family, adiponectin, the second example of the fat hormones, has drawn much attention recently from researchers seeking its biological and clinical implications.¹³

Basic Science of Adiponectin

In the mid-1990s, four independent Japanese or American groups isolated adiponectin (also known as Acrp30/adipoQ/apM1/GBP28) by using diverse techniques.^{14–17} Adiponectin is secreted primarily from mature adipocytes. It is a 30-kDa molecule that contains a 20-residue amino-terminal signal sequence, a variable region, a collagenous domain

and a carboxy-terminal globular domain.¹⁸ Adiponectin circulates in the blood in higher-order structural forms and at a relatively high concentration compared with the other peptide hormones.¹⁷ These multimeric forms identified by different methods (such as gel filtration chromatography, gel electrophoresis or Western blotting) include trimeric (low molecular weight, LMW), hexameric (middle molecular weight, MMW), and even high molecular weight (HMW) complexes.^{17,19} Although some studies have proposed that the ratio of the HMW to other forms may serve as a better indicator of metabolic disorders,^{19–22} the majority of studies that have linked adiponectin with metabolic diseases have used assays for total adiponectin.

One of the effects of adiponectin on energy balance is mediated by an increase in β -oxidation of fatty acids in skeletal muscles.²³ In mice, administration of the protease-generated globular head domain of adiponectin decreases dramatically the level of plasma free fatty acids.²³ It has also been demonstrated in mice that injection of adiponectin can reduce plasma glucose levels and improve insulin sensitivity.^{24,25} The repression of hepatic glucose output is accomplished through lowering the mRNA expression of hepatic gluconeogenic enzymes, such as phosphoenolpyruvate carboxykinase and glucose-6-phosphatase.²⁶ Besides, studies that have utilized adiponectin knockout mice have shown that these mice are inclined to diet-induced insulin resistance and glucose intolerance.^{27,28} In *ob/ob* mice, transgenic adiponectin keeps the animals from developing insulin resistance and hyperglycemia.²⁹

Substantial evidence in animal studies has shown that adiponectin may also exert advantageous effects on the cardiovascular system. In adiponectin knockout mice, hypertension induced by treatment with high-salt diet can be alleviated by adenovirus-delivered adiponectin.³⁰ In adiponectin-deficient mice, pressure overload induces concentric cardiac hypertrophy and mortality.³¹ Moreover, adiponectin supplementation reduces lipid accumulation on the vascular wall,^{29,32} attenuates neointimal proliferation³³

and thrombus formation,³⁴ and improves angiogenic repair.³⁵

In 2003, Yamauchi et al identified adiponectin receptors 1 and 2 (AdipoR1 and AdipoR2).³⁶ These receptors, predicted to contain seven transmembrane domains, are mainly found in skeletal muscle, liver, and pancreatic β cells. AdipoR1 is expressed extensively in skeletal muscle, while AdipoR2 is distributed mostly in liver. Additionally, functional differences between AdipoR1 and AdipoR2 have been demonstrated.³⁷ While AdipoR1 is involved in the AMP-dependent protein kinase (AMPK) signaling pathway, AdipoR2 is associated more with the peroxisome proliferator activator receptor (PPAR)- α pathway.³⁷ Besides the above-mentioned pathways, adiponectin stimulates various signaling pathways. For example, adiponectin activates p38 mitogen-activated protein kinase (MAPK) and c-jun N-terminal kinase (JNK) in osteoblasts, which enhances cell proliferation and differentiation.³⁸ In human umbilical vein endothelial cells, adiponectin inhibits the activation of I κ B kinase (IKK) induced by tumor necrosis factor (TNF)- α or high glucose, which supports its anti-inflammatory role.³⁹ In vascular endothelial cells, adiponectin stimulates production of nitric oxide through phosphatidylinositol-3-kinase-dependent pathways.⁴⁰ The regulation of adiponectin expression and its post-receptor signaling has been discussed in some recent review articles.^{13,41}

Implications of Plasma Adiponectin in Human Cardiometabolic Disorders

Assays for blood adiponectin

For determination of adiponectin levels in peripheral circulation, many immunometric systems, such as radioimmunoassay (RIA) and ELISA, have been developed and have been available commercially for many years. The pioneer development of ELISA was achieved by Arita et al from Matsuzawa's laboratory and has had many applications in clinical research.⁴² For ELISA measurement of adiponectin, the adiponectin multimers

in the biological samples are converted to monomers or dimers by treatment with SDS at high temperature or at pH 3.0–3.5.^{42,43} The calibration curve is obtained by using recombinant adiponectin or dimeric adiponectin as a calibration standard.^{42,43} If a particular molecular weight form is of special interest, two kinds of proteinase can be used to digest selectively the adiponectin multimers.⁴³ Protease A (from *Aspergillus oryzae*), which digests LMW adiponectin can be utilized in the measurement of MMW and HMW forms. In contrast, protease K (from *Tritirachium album*), which digests LMW and MMW adiponectin can be applied to the evaluation of the HMW form. Thus, by combination, the concentration levels of each form may be obtained. For interested readers, a more comprehensive discussion can be found in some recent papers.^{19,43–47} In the following discussion, whenever adiponectin is mentioned, total adiponectin is implied unless indicated otherwise. Previous studies have shown different results with regard to whether HMW adiponectin complexes are the main component related to its biological effects. This will be reviewed in this article. Furthermore, since the amount of the HMW form is so closely related to the total adiponectin level, some authors have reported that measurement of the HMW complexes does not provide additional information.⁴⁸

The association between plasma adiponectin levels and various diseases has been demonstrated in numerous human studies.^{49–51} In this review, we will limit our discussion to human metabolic and cardiovascular disorders. Human genetic studies of adiponectin and its receptors have also revealed important implications of adiponectin in human diseases.^{52–54}

Plasma adiponectin and obesity

Obesity provided the first link between human disease and plasma adiponectin levels.⁴² In the very first report of ELISA for plasma adiponectin, Arita et al demonstrated the paradoxical association between body mass index (BMI) and plasma adiponectin levels.⁴² They also showed that females in general had higher plasma adiponectin than

males. In addition, body fat distribution was found to be influential. Intra-abdominal fat mass is more related to adiponectin concentration than subcutaneous fat is.⁵⁵⁻⁵⁷ Although adiponectin has been related closely to obesity in cross-sectional studies, lower plasma adiponectin is unlikely to be the cause of obesity. On the contrary, hypoadiponectinemia is probably the result of obesity. In fact, in a prospective human study, low plasma adiponectin was not shown to predict future body weight gain.⁵⁸ Recently, we have reviewed extensively this issue.¹³

Lifestyle modification to reduce body weight has been shown to increase plasma adiponectin levels in type 2 diabetes mellitus (T2DM) as well as in non-diabetic subjects.⁵⁹ Exercise itself, without significant weight reduction, may not have a major influence on plasma adiponectin levels.^{60,61} Anti-obesity pharmacotherapy may also increase plasma adiponectin levels. Modest weight reduction supplemented with orlistat, a pancreatic lipase inhibitor, increases plasma adiponectin.^{62,63} Sibutramine, a neurotransmitter reuptake inhibitor, reduces weight with or without increasing plasma adiponectin.^{62,64,65} Rimonabant, an endocannabinoid receptor blocker, reduces weight, improves metabolic profiles and increases plasma adiponectin.⁶⁶ Among severely obese subjects, anti-obesity surgery results in significant weight reduction and metabolic benefits, accompanied by increased plasma adiponectin levels.^{67,68} In contrast, contradictory results regarding the effects of liposuction on obesity-related metabolic abnormalities and plasma adiponectin levels have been reported.^{69,70} In summary, we can conclude that any clinical measure that gives rise to significant weight reduction and improvement in insulin sensitivity can, without doubt, increase the blood concentration of adiponectin.

In more recent studies, the relationship between obesity and different isoforms of adiponectin has been addressed. It appears that total and HMW adiponectin isoforms are more related to obesity and visceral fat.^{71,72} However, some authors have not observed the significance of HMW adiponectin in obesity.⁷³ Weight reduction also appears to

increase total adiponectin, HMW adiponectin and the ratio of HMW to total adiponectin.^{74,75}

Plasma adiponectin and diabetes mellitus

The circulation levels of adiponectin are lower in T2DM,^{59,76} and in insulin-resistant states.^{76,77} The Pima Indians of Arizona, USA, are a special population that is prone to obesity and T2DM. A longitudinal study has revealed that subjects with high adiponectin concentrations are less likely to develop T2DM.⁷⁸ Research on Japanese and other ethnic groups has implied that lower adiponectin levels are associated with increased future risk of developing insulin resistance, impaired glucose metabolism and diabetes.⁷⁹⁻⁸⁵ Reduction of HMW adiponectin isoforms is associated with the risk of insulin resistance and T2DM.⁸⁶⁻⁸⁹

In contrast to that in T2DM, the plasma concentration of adiponectin tends to be higher in type 1 diabetes mellitus (T1DM).⁹⁰⁻⁹³ Even HMW adiponectin is increased in T1DM.⁹⁴ However, in one study, the plasma adiponectin levels did not predict the occurrence of T1DM in autoantibody-positive relatives.⁹⁵ The controversy regarding whether the increased plasma adiponectin concentration is caused by or related to co-existing microvascular complications, especially nephropathy, remains unresolved.^{92,94,96-99}

Lifestyle modification with significant weight reduction has been shown to enhance plasma adiponectin concentration in T2DM, as discussed above.^{59,100} Treatment of T2DM with oral anti-diabetic agents may also alter blood concentrations of adiponectin. In general, insulin secretagogues, including sulfonylureas and meglitinides,^{101,102} do not clearly increase plasma adiponectin levels, except for glimepiride and repaglinide.^{103,104} The rare exception of glimepiride might be explained by its long-claimed insulin-mimetic or insulin-sensitizing effects.¹⁰⁵ However, how repaglinide increases plasma adiponectin remains unknown.¹⁰⁶ Metformin does not appear to have a major effect on plasma adiponectin concentration in T2DM or in polycystic ovary syndrome, even though some studies have shown a significant increase after treatment.^{102,107,108} The α -glucosidase inhibitor

acarbose induced a small but significant rise in serum adiponectin in only one study.¹⁰⁹ In contrast, thiazolidinediones (TZDs), such as rosiglitazone and pioglitazone, increased plasma adiponectin in many human studies.^{101,110} This is related to their insulin-sensitizing effects via PPAR γ 2 activation in the adipose tissue. It has been shown that some of the benefits of TZDs may be exerted mainly through improving the levels of HMW adiponectin complexes.^{111,112} So far, there have been no human studies of dipeptidyl protease-4 inhibitor or glucagon-like peptide-1 (GLP-1)-based therapy in relation to adiponectin. However, we believe, with its effects on body weight, GLP-1-based therapy should more or less increase plasma adiponectin levels. A few studies of insulin treatment have indicated that there might be no significant change in blood adiponectin with chronic insulin treatment.^{113–115} Some other studies have found acute suppressive effects of insulin on adiponectin levels in human studies.^{116–119} However, others have found no effect on plasma adiponectin levels in patients with acute hyperinsulinemia.¹²⁰

Plasma adiponectin and dyslipidemia

Low blood levels of adiponectin are also related to dyslipidemia.^{77,121,122} A positive correlation has been found between adiponectin levels and high-density lipoprotein (HDL)-cholesterol, while an inverse relationship with triglycerides and apolipoprotein (Apo) B has been demonstrated.^{77,121–124} Furthermore, HMW adiponectin was correlated with the decrease in large triglyceride-rich very-low-density lipoprotein (VLDL) and in small low-density lipoprotein (LDL), but with the increase in large buoyant LDL particles, accompanied by increased LDL particle size, and the increase in large HDL and HDL particle size.¹²⁵

Statins are the mainstream pharmacotherapy for hypercholesterolemia, specifically high LDL-cholesterol. Treatment with statins in humans does not significantly alter plasma adiponectin levels in general.^{126–129} The intestinal cholesterol transporter inhibitor ezetimibe is available in clinical practice for treating hypercholesterolemia.

Treatment with ezetimibe also does not alter plasma adiponectin levels.¹³⁰ Fibrates are used to treat hypertriglyceridemia. The effects of fibrates on plasma adiponectin are inconsistent. Some studies have shown an adiponectin-raising effect,^{131,132} while others have not found any effect.^{133,134} One report has suggested that fibrates do not increase total plasma adiponectin, but rather increase the HMW portion.¹³⁵ Nicotinic acid lowers triglycerides, LDL-cholesterol and raises HDL-cholesterol. Extended-released nicotinic acid increases plasma adiponectin, preferentially the HMW forms.^{136–138} Chronic treatment with acipimox, a lipolysis inhibitor similar to nicotinic acid, does not alter plasma adiponectin level.^{139,140} However, the acute effects of acipimox are inconsistent. One study has shown reduced plasma adiponectin levels,¹⁴¹ but another has shown no change in plasma adiponectin concentration after acute administration.¹¹⁷

Plasma adiponectin and hypertension

The inverse relationship between adiponectin level and blood pressure has been investigated in many previous studies,^{142–145} and re-enforced recently by a 5-year prospective study.¹⁴⁶ Chow et al found that hypoadiponectinemia can serve as a powerful predictor for future development of hypertension, even after adjustment for well-known risk factors, including sex, age and BMI.¹⁴⁶

Antihypertensive medication may modulate the plasma levels of adiponectin. Thiazide diuretics seem to reduce plasma adiponectin levels.¹⁴⁷ Another diuretic, indapamide, also reduces plasma adiponectin concentration.¹⁴⁸ Beta-adrenergic blockers reduce or have no effect on plasma adiponectin levels.^{147–149} Calcium channel blockers increase or have no effect on plasma adiponectin concentrations.^{147,149–151} Blocking the renin-angiotensin-aldosterone system with angiotensin-converting enzyme inhibitors or angiotensin receptor 1 blockers consistently increase plasma adiponectin levels.^{126,147,149,152–154} Spironolactone has also been reported to increase plasma adiponectin in type 2 diabetic patients with nephropathy.¹⁵⁵ No human studies of eplerenone or

aliskiren have been published. The α -adrenergic blocker doxazosin has also been shown to increase plasma adiponectin levels.¹⁴⁹ Overall, the adiponectin-raising effects of antihypertensive agents appear to parallel their effects on insulin sensitivity.^{147,149} Among these antihypertensives, so far, only losartan has been shown to increase total and HMW adiponectin.¹⁵⁶

Hypoadiponectinemia is related closely to obesity, T2DM, dyslipidemia and hypertension. These metabolic abnormalities are not only the key components of the metabolic syndrome, but also the major risk factors for cardiovascular diseases.⁷⁷ Therefore, we will next review the relationship between blood adiponectin and coronary, cerebral and peripheral artery diseases, congestive heart failure (CHF) and cardiac arrhythmia.

Plasma adiponectin and coronary artery disease (CAD)

Plasma adiponectin was first shown to be lower in patients with CAD by Ouchi et al in 1999.¹⁵⁷ The association between low plasma adiponectin and CAD, disease extent or outcomes has been demonstrated not only in cross-sectional studies,^{157–161} but also in prospective longitudinal follow-up studies.^{162–166} For example, in a follow-up study of male health professionals, the subjects with higher adiponectin levels had a significantly reduced risk of myocardial infarction, even after correction for risk factors, such as LDL and HDL levels, BMI, history of diabetes and hypertension and high-sensitivity C-reactive protein at baseline.¹⁶⁷ Some other studies, however, have not found a predictive value for future risk of CAD.^{168,169} A meta-analysis has suggested a significant association, but the effect was relatively moderate compared with that in previous studies.¹⁷⁰

HMW adiponectin is lower in men with CAD, but not in women.¹⁷¹ In a follow-up study, HMW adiponectin has been shown to predict cardiovascular events in men.¹⁷² HMW adiponectin is also associated with extent of CAD in men.¹⁷³ However, negative results have also been reported.^{174–177} Whether gender is an issue or not

in this regard remains to be clarified. Furthermore, whether adiponectin plays a specific role at certain specific stages of CAD pathogenesis needs to be investigated more thoroughly.

Plasma adiponectin and stroke and peripheral artery disease (PAD)

Lower blood adiponectin levels have been reported to be associated with increased risk of ischemic stroke and increased mortality after first stroke event.^{178,179} On the other hand, some other studies have not confirmed the value of adiponectin in relation to cerebrovascular events.^{180–182} Plasma adiponectin has also been shown to be inversely related to PAD, estimated by ankle-brachial index in hemodialysis patients.¹⁸³ So far, no study has addressed the role of HMW adiponectin isoforms in cerebrovascular disease or PAD. Furthermore, no human investigation of the effects of pentoxifylline or cilostazol, which are commonly used in treating PAD in clinical practice, on plasma adiponectin levels has been reported. However, a novel selective serotonin 2A receptor antagonist, sarpogrelate, has been investigated in non-diabetic and non-medicated diabetic patients with PAD.¹⁸⁴ It was shown to enhance insulin sensitivity and increase plasma adiponectin.

Plasma adiponectin and CHF and cardiac arrhythmia

The first human study regarding the role of adiponectin in CHF was surprising.¹⁸⁵ It showed that high blood adiponectin was a predictor for mortality in CHF, independent of other risk factors. It was suggested that high adiponectin in CHF was a sign of physical wasting. Other studies have also demonstrated that higher adiponectin is an independent predictor for mortality and/or hospitalization in patients with CHF.^{186–189} In cross-sectional studies, patients with CHF had higher plasma adiponectin levels and decreased lean body mass.^{188,190,191} However, in a longitudinal follow-up study, in which insulin resistance was reported to be an independent risk factor for CHF, plasma adiponectin was not a predictor for

CHF.¹⁹² In one study that has surveyed the role of HMW adiponectin, it was concluded that total, rather than HMW adiponectin was valuable in predicting mortality in CHF.¹⁹³ In many studies, plasma adiponectin level appears to parallel the level of natriuretic peptide.^{185,186,188,189,191,194,195} Moreover, infusion of carperitide (a natriuretic peptide) in CHF patients increased their plasma total and HMW adiponectin levels.¹⁹⁶

Cardiac arrhythmia is a common condition in patients with cardiovascular diseases. It has been reported that plasma adiponectin levels are higher in chronic atrial fibrillation (AF) than in paroxysmal AF and controls.¹⁹⁷ The plasma adiponectin levels are correlated with the collagen metabolism marker, carboxy-terminal telopeptide of collagen I, but not with type III pro-collagen-N-peptide.¹⁹⁷ So far, there has been no report of adiponectin with regard to ventricular arrhythmia, syncope, fainting or sudden death.

Conclusion

Substantial evidence has revealed the significance of adiponectin in human health and diseases. The underlying molecular mechanisms have provided us with clues to the intricate relationship between adiponectin and pathogenesis. Alterations in blood adiponectin concentration seem to be a useful biomarker for obesity, DM, dyslipidemia, hypertension, and even subsequent cardiovascular diseases. We summarized the relationship of blood adiponectin with cardiometabolic disorders in this review. The proposed natural history is depicted in the Figure. Hypoadiponectinemia may be caused by obesity-induced insulin resistance in the adipose tissue, as previously proposed.¹³ Hypoadiponectinemia may contribute secondarily to insulin resistance in other peripheral tissues, such as liver and skeletal muscle.¹³ As a result, T2DM, dyslipidemia, hypertension, metabolic syndrome and




Figure. Schematic diagram depicting the proposed natural course of blood adiponectin levels in relation to cardiometabolic disorders reviewed in this article. The arrows indicate the proposed time sequence and may not necessarily imply cause–effect relationships. The “+” signs indicate a positive association. The “–” sign indicates a negative association. Hypoadiponectinemia is proposed to be caused by obesity-induced insulin resistance in the adipose tissue.¹³ Hypoadiponectinemia in turn induces insulin resistance in other tissues. As the result of certain microvascular complications, such as nephropathy and CHF, the plasma levels of adiponectin may rise. T2DM = type 2 diabetes mellitus; CAD = coronary artery disease; CVA = cerebrovascular accident; PAD = peripheral artery disease; CHF = congestive heart failure; AF = atrial fibrillation.

accelerated atherosclerosis occur. These conditions may lead to subsequent macro- and microvascular complications, and CHF and arrhythmia may follow. As a result of CHF, AF or nephropathy, compensatory or secondary rise in blood adiponectin may occur later (Figure). The mechanisms of this rise in blood adiponectin levels remain unclear. It may be caused by reduced renal clearance of adiponectin, disease-related weight loss, or increased ectopic secretion of adiponectin from the myocardium. If the rise in blood adiponectin is proven to be beneficial, we do not have to wait until late clinical events occur. Non-pharmacological and pharmacological modulations of blood adiponectin levels may be promising for ameliorating the related disorders in the earlier stages. Through more comprehensive studies, we believe that the true clinical value and the mystery of the pathogenic mechanisms of adiponectin in human diseases will be unraveled in the near future.

Acknowledgments

Dr Lin-Chau Chang is grateful to the National Taiwan University for her postdoctoral fellowship from "Aim for Top University Project". Dr Wei-Shiung Yang is in part supported by grant 97-2314-B-002-050-MY3 from the National Science Council of Taiwan for the research on adiponectin.

References

1. Grundy SM. Metabolic syndrome pandemic. *Arterioscler Thromb Vasc Biol* 2008;28:629–36.
2. Hampton T. Scientists study fat as endocrine organ. *JAMA* 2006;296:1573–5.
3. Ingalls AM, Dickie MM, Snell GD. Obese, a new mutation in the house mouse. *J Hered* 1950;41:317–8.
4. Hummel KP, Dickie MM, Coleman DL. Diabetes, a new mutation in the mouse. *Science* 1966;153:1127–8.
5. Coleman DL. Obese and diabetes: two mutant genes causing diabetes–obesity syndromes in mice. *Diabetologia* 1978;14:141–8.
6. Coleman DL. Effects of parabiosis of obese with diabetes and normal mice. *Diabetologia* 1973;9:294–8.
7. Coleman DL, Hummel KP. Effects of parabiosis of normal with genetically diabetic mice. *Am J Physiol* 1969;217:1298–304.
8. Zhang Y, Proenca R, Maffei M, et al. Positional cloning of the mouse obese gene and its human homologue. *Nature* 1994;372:425–32.
9. Halaas JL, Gajiwala KS, Maffei M, et al. Weight-reducing effects of the plasma protein encoded by the obese gene. *Science* 1995;269:543–6.
10. Tartaglia LA, Dembski M, Weng X, et al. Identification and expression cloning of a leptin receptor, ob-r. *Cell* 1995;83:1263–71.
11. Myers MG, Cowley MA, Munzberg H. Mechanisms of leptin action and leptin resistance. *Annu Rev Physiol* 2008;70:537–56.
12. Leininger GM, Myers MG Jr. Lrb signals act within a distributed network of leptin-responsive neurones to mediate leptin action. *Acta Physiol (Oxf)* 2008;192:49–59.
13. Lu JY, Huang KC, Chang LC, et al. Adiponectin: a biomarker of obesity-induced insulin resistance in adipose tissue and beyond. *J Biomed Sci* 2008;15:565–76.
14. Scherer PE, Williams S, Fogliano M, et al. A novel serum protein similar to c1q, produced exclusively in adipocytes. *J Biol Chem* 1995;270:26746–9.
15. Maeda K, Okubo K, Shimomura I, et al. Cdn cloning and expression of a novel adipose specific collagen-like factor, apm1 (adipose most abundant gene transcript 1). *Biochem Biophys Res Commun* 1996;221:286–9.
16. Hu E, Liang P, Spiegelman BM. Adipoq is a novel adipose-specific gene dysregulated in obesity. *J Biol Chem* 1996;271:10697–703.
17. Nakano Y, Tobe T, Choi-Miura NH, et al. Isolation and characterization of gbp28, a novel gelatin-binding protein purified from human plasma. *J Biochem (Tokyo)* 1996;120:803–12.
18. Pajvani UB, Du X, Combs TP, et al. Structure–function studies of the adipocyte-secreted hormone acrp30/adiponectin. Implications for metabolic regulation and bioactivity. *J Biol Chem* 2003;278:9073–85.
19. Magkos F, Sidossis LS. Recent advances in the measurement of adiponectin isoform distribution. *Curr Opin Clin Nutr Metab Care* 2007;10:571–5.
20. Waki H, Yamauchi T, Kamon J, et al. Impaired multimerization of human adiponectin mutants associated with diabetes. Molecular structure and multimer formation of adiponectin. *J Biol Chem* 2003;278:40352–63.
21. Kobayashi H, Ouchi N, Kihara S, et al. Selective suppression of endothelial cell apoptosis by the high molecular weight form of adiponectin. *Circ Res* 2004;94:e27–31.
22. Pajvani UB, Hawkins M, Combs TP, et al. Complex distribution, not absolute amount of adiponectin, correlates with thiazolidinedione-mediated improvement in insulin sensitivity. *J Biol Chem* 2004;279:12152–62.
23. Fruebis J, Tsoo TS, Javorschi S, et al. Proteolytic cleavage product of 30-kda adipocyte complement-related protein

- increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci USA* 2001;98:2005–10.
24. Berg AH, Combs TP, Du X, et al. The adipocyte-secreted protein acrp30 enhances hepatic insulin action. *Nat Med* 2001;7:947–53.
 25. Yamauchi T, Kamon J, Waki H, et al. The fat-derived hormone adiponectin reverses insulin resistance associated with both lipodystrophy and obesity. *Nat Med* 2001;7:941–6.
 26. Combs TP, Berg AH, Obici S, et al. Endogenous glucose production is inhibited by the adipose-derived protein acrp30. *J Clin Invest* 2001;108:1875–81.
 27. Kubota N, Terauchi Y, Yamauchi T, et al. Disruption of adiponectin causes insulin resistance and neointimal formation. *J Biol Chem* 2002;277:25863–6.
 28. Maeda N, Shimomura I, Kishida K, et al. Diet-induced insulin resistance in mice lacking adiponectin/acrp30. *Nat Med* 2002;8:731–7.
 29. Yamauchi T, Kamon J, Waki H, et al. Globular adiponectin protected ob/ob mice from diabetes and apoE-deficient mice from atherosclerosis. *J Biol Chem* 2003;278:2461–8.
 30. Ohashi K, Kihara S, Ouchi N, et al. Adiponectin replenishment ameliorates obesity-related hypertension. *Hypertension* 2006;47:1108–16.
 31. Shibata R, Ouchi N, Ito M, et al. Adiponectin-mediated modulation of hypertrophic signals in the heart. *Nat Med* 2004;10:1384–9.
 32. Okamoto Y, Kihara S, Ouchi N, et al. Adiponectin reduces atherosclerosis in apolipoprotein e-deficient mice. *Circulation* 2002;106:2767–70.
 33. Matsuda M, Shimomura I, Sata M, et al. Role of adiponectin in preventing vascular stenosis. The missing link of adipo-vascular axis. *J Biol Chem* 2002;277:37487–91.
 34. Kato H, Kashiwagi H, Shiraga M, et al. Adiponectin acts as an endogenous antithrombotic factor. *Arterioscler Thromb Vasc Biol* 2006;26:224–30.
 35. Shibata R, Ouchi N, Kihara S, et al. Adiponectin stimulates angiogenesis in response to tissue ischemia through stimulation of amp-activated protein kinase signaling. *J Biol Chem* 2004;279:28670–4.
 36. Yamauchi T, Kamon J, Ito Y, et al. Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 2003;423:762–9.
 37. Yamauchi T, Nio Y, Maki T, et al. Targeted disruption of adipor1 and adipor2 causes abrogation of adiponectin binding and metabolic actions. *Nat Med* 2007;13:332–9.
 38. Luo XH, Guo LJ, Yuan LQ, et al. Adiponectin stimulates human osteoblasts proliferation and differentiation via the MAPK signaling pathway. *Exp Cell Res* 2005;309:99–109.
 39. Wu X, Mahadev K, Fuchsel L, et al. Adiponectin suppresses I κ B kinase activation induced by tumor necrosis factor- α or high glucose in endothelial cells: role of camp and amp kinase signaling. *Am J Physiol Endocrinol Metab* 2007;293:E1836–44.
 40. Chen H, Montagnani M, Funahashi T, et al. Adiponectin stimulates production of nitric oxide in vascular endothelial cells. *J Biol Chem* 2003;278:45021–6.
 41. Kadowaki T, Yamauchi T, Kubota N. The physiological and pathophysiological role of adiponectin and adiponectin receptors in the peripheral tissues and CNS. *FEBS Lett* 2008;582:74–80.
 42. Arita Y, Kihara S, Ouchi N, et al. Paradoxical decrease of an adipose-specific protein, adiponectin, in obesity. *Biochem Biophys Res Commun* 1999;257:79–83.
 43. Ebinuma H, Miyazaki O, Yago H, et al. A novel ELISA system for selective measurement of human adiponectin multimers by using proteases. *Clin Chim Acta* 2006;372:47–53.
 44. Nakano Y, Tajima S, Yoshimi A, et al. A novel enzyme-linked immunosorbent assay specific for high-molecular-weight adiponectin. *J Lipid Res* 2006;47:1572–82.
 45. Ebinuma H, Miida T, Yamauchi T, et al. Improved ELISA for selective measurement of adiponectin multimers and identification of adiponectin in human cerebrospinal fluid. *Clin Chem* 2007;53:1541–4.
 46. Tanita T, Miyakoshi H, Nakano Y. Performance of ELISA for specific measurement of high-molecular-weight (HMW) adiponectin. *J Immunol Methods* 2008;333:139–46.
 47. Sinha MK, Songer T, Xiao Q, et al. Analytical validation and biological evaluation of a high molecular-weight adiponectin ELISA. *Clin Chem* 2007;53:2144–51.
 48. Glinborg D, Frystyk J, Hojlund K, et al. Total and high molecular weight (HMW) adiponectin levels and measures of glucose and lipid metabolism following pioglitazone treatment in a randomized placebo-controlled study in polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2008;68:165–74.
 49. Rogers CQ, Ajmo JM, You M. Adiponectin and alcoholic fatty liver disease. *IUBMB Life* 2008;60:790–7.
 50. Gualillo O, Gonzalez-Juanatey JR, Lago F. The emerging role of adipokines as mediators of cardiovascular function: physiologic and clinical perspectives. *Trends Cardiovasc Med* 2007;17:275–83.
 51. Kelesidis I, Kelesidis T, Mantzoros CS. Adiponectin and cancer: a systematic review. *Br J Cancer* 2006;94:1221–5.
 52. Yang WS, Chuang LM. Human genetics of adiponectin in the metabolic syndrome. *J Mol Med* 2006;84:112–21.
 53. Menzaghi C, Trischitta V, Doria A. Genetic influences of adiponectin on insulin resistance, type 2 diabetes, and cardiovascular disease. *Diabetes* 2007;56:1198–209.
 54. Crimmins NA, Woo JG, Kaushal RD, et al. Adiponectin receptor 1 variants associated with lower insulin resistance in African Americans. *Obesity* 2007;15:1903–7.
 55. Cnop M, Havel PJ, Utzschneider KM, et al. Relationship of adiponectin to body fat distribution, insulin sensitivity and plasma lipoproteins: evidence for independent roles of age and sex. *Diabetologia* 2003;46:459–69.
 56. Yatagai T, Nagasaka S, Taniguchi A, et al. Hypoadiponectinemia is associated with visceral fat accumulation and insulin resistance in Japanese men with type 2 diabetes mellitus. *Metabolism* 2003;52:1274–8.

57. Staiger H, Tschritter O, Machann J, et al. Relationship of serum adiponectin and leptin concentrations with body fat distribution in humans. *Obes Res* 2003;11:368–72.
58. Vozarova B, Stefan N, Lindsay RS, et al. Low plasma adiponectin concentrations do not predict weight gain in humans. *Diabetes* 2002;51:2964–7.
59. Hotta K, Funahashi T, Arita Y, et al. Plasma concentrations of a novel, adipose-specific protein, adiponectin, in type 2 diabetic patients. *Arterioscler Thromb Vasc Biol* 2000;20:1595–9.
60. Hulver MW, Zheng D, Tanner CJ, et al. Adiponectin is not altered with exercise training despite enhanced insulin action. *Am J Physiol Endocrinol Metab* 2002;283:E861–5.
61. Yatagai T, Nishida Y, Nagasaka S, et al. Relationship between exercise training-induced increase in insulin sensitivity and adiponectinemia in healthy men. *Endocr J* 2003;50:233–8.
62. Valsamakis G, McTernan PG, Chetty R, et al. Modest weight loss and reduction in waist circumference after medical treatment are associated with favorable changes in serum adipocytokines. *Metabolism* 2004;53:430–4.
63. Hsieh CJ, Wang PW, Liu RT, et al. Orlistat for obesity: benefits beyond weight loss. *Diabetes Res Clin Pract* 2005;67:78–83.
64. Kim DM, Yoon SJ, Ahn CW, et al. Sibutramine improves fat distribution and insulin resistance, and increases serum adiponectin levels in Korean obese nondiabetic premenopausal women. *Diabetes Res Clin Pract* 2004;66(Suppl 1):S139–44.
65. Hung YJ, Chen YC, Pei D, et al. Sibutramine improves insulin sensitivity without alteration of serum adiponectin in obese subjects with type 2 diabetes. *Diabet Med* 2005;22:1024–30.
66. Despres JP, Golay A, Sjostrom L. Effects of rimonabant on metabolic risk factors in overweight patients with dyslipidemia. *N Engl J Med* 2005;353:2121–34.
67. Calvani M, Scarfone A, Granato L, et al. Restoration of adiponectin pulsatility in severely obese subjects after weight loss. *Diabetes* 2004;53:939–47.
68. Yang WS, Lee WJ, Funahashi T, et al. Weight reduction increases plasma levels of an adipose-derived anti-inflammatory protein, adiponectin. *J Clin Endocrinol Metab* 2001;86:3815–9.
69. Klein S, Fontana L, Young VL, et al. Absence of an effect of liposuction on insulin action and risk factors for coronary heart disease. *N Engl J Med* 2004;350:2549–57.
70. Giugliano G, Nicoletti G, Grella E, et al. Effect of liposuction on insulin resistance and vascular inflammatory markers in obese women. *Br J Plast Surg* 2004;57:190–4.
71. Nakatani H, Hirose H, Yamamoto Y, et al. Significance of leptin and high-molecular weight adiponectin in the general population of Japanese male adolescents. *Metabolism* 2008;57:157–62.
72. Kaser S, Tatarczyk T, Stadlmayr A, et al. Effect of obesity and insulin sensitivity on adiponectin isoform distribution. *Eur J Clin Invest* 2008;38:827–34.
73. Schober F, Neumeier M, Weigert J, et al. Low molecular weight adiponectin negatively correlates with the waist circumference and monocytic IL-6 release. *Biochem Biophys Res Commun* 2007;361:968–73.
74. Bobbert T, Rochlitz H, Wegewitz U, et al. Changes of adiponectin oligomer composition by moderate weight reduction. *Diabetes* 2005;54:2712–9.
75. Engl J, Bobbert T, Ciardi C, et al. Effects of pronounced weight loss on adiponectin oligomer composition and metabolic parameters. *Obesity (Silver Spring)* 2007;15:1172–8.
76. Weyer C, Funahashi T, Tanaka S, et al. Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J Clin Endocrinol Metab* 2001;86:1930–5.
77. Yang WS, Lee WJ, Funahashi T, et al. Plasma adiponectin levels in overweight and obese Asians. *Obes Res* 2002;10:1104–10.
78. Lindsay RS, Funahashi T, Hanson RL, et al. Adiponectin and development of type 2 diabetes in the Pima Indian population. *Lancet* 2002;360:57–8.
79. Daimon M, Oizumi T, Saitoh T, et al. Decreased serum levels of adiponectin are a risk factor for the progression to type 2 diabetes in the Japanese population: the Funagata study. *Diabetes Care* 2003;26:2015–20.
80. Snehathatha C, Mukesh B, Simon M, et al. Plasma adiponectin is an independent predictor of type 2 diabetes in Asian Indians. *Diabetes Care* 2003;26:3226–9.
81. Stefan N, Vozarova B, Funahashi T, et al. Plasma adiponectin concentration is associated with skeletal muscle insulin receptor tyrosine phosphorylation, and low plasma concentration precedes a decrease in whole-body insulin sensitivity in humans. *Diabetes* 2002;51:1884–8.
82. Yamamoto Y, Hirose H, Saito I, et al. Adiponectin, an adipocyte-derived protein, predicts future insulin resistance: two-year follow-up study in Japanese population. *J Clin Endocrinol Metab* 2004;89:87–90.
83. Duncan BB, Schmidt MI, Pankow JS, et al. Adiponectin and the development of type 2 diabetes: the atherosclerosis risk in communities study. *Diabetes* 2004;53:2473–8.
84. Snijder MB, Heine RJ, Seidell JC, et al. Associations of adiponectin levels with incident impaired glucose metabolism and type 2 diabetes in older men and women: the Hoorn study. *Diabetes Care* 2006;29:2498–503.
85. Mather KJ, Funahashi T, Matsuzawa Y, et al. Adiponectin, change in adiponectin, and progression to diabetes in the diabetes prevention program. *Diabetes* 2008;57:980–6.
86. Hara K, Horikoshi M, Yamauchi T, et al. Measurement of the high-molecular weight form of adiponectin in plasma is useful for the prediction of insulin resistance and metabolic syndrome. *Diabetes Care* 2006;29:1357–62.
87. Nakashima R, Kamei N, Yamane K, et al. Decreased total and high molecular weight adiponectin are independent risk factors for the development of type 2 diabetes in Japanese-Americans. *J Clin Endocrinol Metab* 2006;91:3873–7.

88. Basu R, Pajvani UB, Rizza RA, et al. Selective downregulation of the high-molecular weight form of adiponectin in hyperinsulinemia and in type 2 diabetes. Differential regulation from nondiabetic subjects. *Diabetes* 2007; 56:2174–7.
89. Seino Y, Hirose H, Saito I, et al. High molecular weight multimer form of adiponectin as a useful marker to evaluate insulin resistance and metabolic syndrome in Japanese men. *Metabolism* 2007;56:1493–9.
90. Imagawa A, Funahashi T, Nakamura T, et al. Elevated serum concentration of adipose-derived factor, adiponectin, in patients with type 1 diabetes. *Diabetes Care* 2002;25:1665–6.
91. Celi F, Bini V, Papi F, et al. Circulating adipocytokines in non-diabetic and type 1 diabetic children: relationship to insulin therapy, glycaemic control and pubertal development. *Diabet Med* 2006;23:660–5.
92. Frystyk J, Tarnow L, Hansen TK, et al. Increased serum adiponectin levels in type 1 diabetic patients with microvascular complications. *Diabetologia* 2005;48:1911–8.
93. Galler A, Gelbrich G, Kratzsch J, et al. Elevated serum levels of adiponectin in children, adolescents and young adults with type 1 diabetes and the impact of age, gender, body mass index and metabolic control: a longitudinal study. *Eur J Endocrinol* 2007;157:481–9.
94. Leth H, Andersen KK, Frystyk J, et al. Elevated levels of high-molecular-weight adiponectin in type 1 diabetes. *J Clin Endocrinol Metab* 2008;93:3186–91.
95. Truyen I, De Grijse J, Van Schravendijk C, et al. Adiponectin levels do not predict clinical onset of type 1 diabetes in antibody-positive relatives. *Diabetologia* 2007; 50:2143–6.
96. Schaffler A, Herfarth H, Paul G, et al. Identification of influencing variables on adiponectin serum levels in diabetes mellitus type 1 and type 2. *Exp Clin Endocrinol Diabetes* 2004;112:383–9.
97. Schalkwijk CG, Chaturvedi N, Schram MT, et al. Adiponectin is inversely associated with renal function in type 1 diabetic patients. *J Clin Endocrinol Metab* 2006; 91:129–35.
98. Saraheimo M, Forsblom C, Thorn L, et al. Serum adiponectin and progression of diabetic nephropathy in patients with type 1 diabetes. *Diabetes Care* 2008;31: 1165–9.
99. Barnes MM, Curran-Everett D, Hamman RF, et al. Determinants of adiponectin levels in young people with type 1 diabetes. *Diabet Med* 2008;25:365–9.
100. Balagopal P, George D, Yarandi H, et al. Reversal of obesity-related hypoadiponectinemia by lifestyle intervention: a controlled, randomized study in obese adolescents. *J Clin Endocrinol Metab* 2005;90:6192–7.
101. Yang WS, Jeng CY, Wu TJ, et al. Synthetic peroxisome proliferator-activated receptor-gamma agonist, rosiglitazone, increases plasma levels of adiponectin in type 2 diabetic patients. *Diabetes Care* 2002;25:376–80.
102. Putz DM, Goldner WS, Bar RS, et al. Adiponectin and C-reactive protein in obesity, type 2 diabetes, and monotherapy. *Metabolism* 2004;53:1454–61.
103. Nagasaka S, Taniguchi A, Aiso Y, et al. Effect of glimepiride on serum adiponectin level in subjects with type 2 diabetes. *Diabetes Care* 2003;26:2215–6.
104. Tsunekawa T, Hayashi T, Suzuki Y, et al. Plasma adiponectin plays an important role in improving insulin resistance with glimepiride in elderly type 2 diabetic subjects. *Diabetes Care* 2003;26:285–9.
105. Muller G. The molecular mechanism of the insulin-mimetic/sensitizing activity of the antidiabetic sulfonylurea drug amaryl. *Mol Med* 2000;6:907–33.
106. Lund SS, Tarnow L, Stehouwer CD, et al. Targeting hyperglycaemia with either metformin or repaglinide in non-obese patients with type 2 diabetes: results from a randomized crossover trial. *Diabetes Obes Metab* 2007;9: 394–407.
107. Spranger J, Mohlig M, Wegewitz U, et al. Adiponectin is independently associated with insulin sensitivity in women with polycystic ovary syndrome. *Clin Endocrinol (Oxf)* 2004;61:738–46.
108. Rasouli N, Yao-Borengasser A, Miles LM, et al. Increased plasma adiponectin in response to pioglitazone does not result from increased gene expression. *Am J Physiol Endocrinol Metab* 2006;290:E42–6.
109. Ochiai H, Ooka H, Shida C, et al. Acarbose treatment increases serum total adiponectin levels in patients with type 2 diabetes. *Endocr J* 2008;55:549–56.
110. Hirose H, Kawai T, Yamamoto Y, et al. Effects of pioglitazone on metabolic parameters, body fat distribution, and serum adiponectin levels in Japanese male patients with type 2 diabetes. *Metabolism* 2002;51: 314–7.
111. Qurashi S, Mynarcik DC, McNurlan MA, et al. Importance of the high-molecular-mass isoform of adiponectin in improved insulin sensitivity with rosiglitazone treatment in HIV disease. *Clin Sci (Lond)* 2008;115:197–202.
112. Tonelli J, Li W, Kishore P, et al. Mechanisms of early insulin-sensitizing effects of thiazolidinediones in type 2 diabetes. *Diabetes* 2004;53:1621–9.
113. Pfitzner A, Lorra B, Abdollahnia MR, et al. The switch from sulfonylurea to preprandial short-acting insulin analog substitution has an immediate and comprehensive beta-cell protective effect in patients with type 2 diabetes mellitus. *Diabetes Technol Ther* 2006;8: 375–84.
114. Miyashita Y, Nishimura R, Nemoto M, et al. Prospective randomized study for optimal insulin therapy in type 2 diabetic patients with secondary failure. *Cardiovasc Diabetol* 2008;7:16.
115. Juurinen L, Tiikkainen M, Hakkinen AM, et al. Effects of insulin therapy on liver fat content and hepatic insulin sensitivity in patients with type 2 diabetes. *Am J Physiol Endocrinol Metab* 2007;292:E829–35.

116. Mohlig M, Wegewitz U, Osterhoff M, et al. Insulin decreases human adiponectin plasma levels. *Horm Metab Res* 2002;34:655–8.
117. Dullaart RP, Riemens SC, Meinardi JR, et al. Plasma adiponectin is modestly decreased during 24-hour insulin infusion but not after inhibition of lipolysis by acipimox. *Scand J Clin Lab Invest* 2005;65:523–31.
118. Brame LA, Considine RV, Yamauchi M, et al. Insulin and endothelin in the acute regulation of adiponectin *in vivo* in humans. *Obes Res* 2005;13:582–8.
119. Blumer RM, van der Crabben SN, Stegenga ME, et al. Hyperglycemia prevents the suppressive effect of hyperinsulinemia on plasma adiponectin levels in healthy humans. *Am J Physiol Endocrinol Metab* 2008;295:E613–7.
120. Krusinova E, Klementova M, Kopecky J, et al. Effect of acute hyperinsulinaemia with and without angiotensin II type 1 receptor blockade on resistin and adiponectin concentrations and expressions in healthy subjects. *Eur J Endocrinol* 2007;157:443–9.
121. Matsubara M, Maruoka S, Katayose S. Decreased plasma adiponectin concentrations in women with dyslipidemia. *J Clin Endocrinol Metab* 2002;87:2764–9.
122. Huang KC, Lue BH, Yen RF, et al. Plasma adiponectin levels and metabolic factors in nondiabetic adolescents. *Obes Res* 2004;12:119–24.
123. Schulze MB, Rimm EB, Shai I, et al. Relationship between adiponectin and glycemic control, blood lipids, and inflammatory markers in men with type 2 diabetes. *Diabetes Care* 2004;27:1680–7.
124. Kazumi T, Kawaguchi A, Hirano T, et al. Serum adiponectin is associated with high-density lipoprotein cholesterol, triglycerides, and low-density lipoprotein particle size in young healthy men. *Metabolism* 2004;53:589–93.
125. Lara-Castro C, Luo N, Wallace P, et al. Adiponectin multimeric complexes and the metabolic syndrome trait cluster. *Diabetes* 2006;55:249–59.
126. Koh KK, Quon MJ, Han SH, et al. Additive beneficial effects of losartan combined with simvastatin in the treatment of hypercholesterolemic, hypertensive patients. *Circulation* 2004;110:3687–92.
127. Bayes B, Granada ML, Lauzurica R, et al. Effect of low doses of atorvastatin on adiponectin, glucose homeostasis, and clinical inflammatory markers in kidney transplant recipients. *Transplant Proc* 2005;37:3808–12.
128. Gannage-Yared MH, Azar RR, Amm-Azar M, et al. Pravastatin does not affect insulin sensitivity and adipocytokines levels in healthy nondiabetic patients. *Metabolism* 2005;54:947–51.
129. Sonmez A, Dogru T, Tasci I, et al. The effect of fluvastatin on plasma adiponectin levels in dyslipidaemia. *Clin Endocrinol (Oxf)* 2006;64:567–72.
130. Gouni-Berthold I, Berthold HK, Chamberland JP, et al. Short-term treatment with ezetimibe, simvastatin or their combination does not alter circulating adiponectin, resistin or leptin levels in healthy men. *Clin Endocrinol (Oxf)* 2008;68:536–41.
131. Koh KK, Quon MJ, Han SH, et al. Additive beneficial effects of fenofibrate combined with candesartan in the treatment of hypertriglyceridemic hypertensive patients. *Diabetes Care* 2006;29:195–201.
132. Rosenson RS. Effect of fenofibrate on adiponectin and inflammatory biomarkers in metabolic syndrome patients. *Obesity (Silver Spring)* 2009;17:504–9.
133. Asano A, Kobayashi J, Murase Y, et al. Effects of fenofibrate therapy on plasma ubiquinol-10 and ubiquinone-10 levels in Japanese patients with hyperlipidemia and type 2 diabetes mellitus. *Pharmacotherapy* 2006;26:447–51.
134. Otto C, Otto B, Frost RJ, et al. Short-term therapy with atorvastatin or fenofibrate does not affect plasma ghrelin, resistin or adiponectin levels in type 2 diabetic patients with mixed hyperlipoproteinaemia. *Acta Diabetol* 2007;44:65–8.
135. Oki K, Koide J, Nakanishi S, et al. Fenofibrate increases high molecular weight adiponectin in subjects with hypertriglyceridemia. *Endocr J* 2007;54:431–5.
136. Westphal S, Borucki K, Taneva E, et al. Extended-release niacin raises adiponectin and leptin. *Atherosclerosis* 2007;193:361–5.
137. Plaisance EP, Grandjean PW, Brunson BL, et al. Increased total and high-molecular weight adiponectin after extended-release niacin. *Metabolism* 2008;57:404–9.
138. Westphal S, Luley C. Preferential increase in high-molecular weight adiponectin after niacin. *Atherosclerosis* 2008;198:179–83.
139. Bajaj M, Suraamornkul S, Kashyap S, et al. Sustained reduction in plasma free fatty acid concentration improves insulin action without altering plasma adipocytokine levels in subjects with strong family history of type 2 diabetes. *J Clin Endocrinol Metab* 2004;89:4649–55.
140. Bajaj M, Suraamornkul S, Romanelli A, et al. Effect of a sustained reduction in plasma free fatty acid concentration on intramuscular long-chain fatty acyl-CoAs and insulin action in type 2 diabetic patients. *Diabetes* 2005;54:3148–53.
141. Bernstein EL, Koutkia P, Ljungquist K, et al. Acute regulation of adiponectin by free fatty acids. *Metabolism* 2004;53:790–3.
142. Kazumi T, Kawaguchi A, Sakai K, et al. Young men with high-normal blood pressure have lower serum adiponectin, smaller LDL size, and higher elevated heart rate than those with optimal blood pressure. *Diabetes Care* 2002;25:971–6.
143. Dzielinska Z, Januszewicz A, Wiecek A, et al. Decreased plasma concentration of a novel anti-inflammatory protein—adiponectin—in hypertensive men with coronary artery disease. *Thromb Res* 2003;110:365–9.

144. Huang KC, Chen CL, Chuang LM, et al. Plasma adiponectin levels and blood pressures in nondiabetic adolescent females. *J Clin Endocrinol Metab* 2003;88: 4130–4.
145. Adamczak M, Wiecek A, Funahashi T, et al. Decreased plasma adiponectin concentration in patients with essential hypertension. *Am J Hypertens* 2003;16:72–5.
146. Chow WS, Cheung BM, Tso AW, et al. Hypoadiponectinemia as a predictor for the development of hypertension: a 5-year prospective study. *Hypertension* 2007;49:1455–61.
147. Koh KK, Quon MJ, Han SH, et al. Distinct vascular and metabolic effects of different classes of anti-hypertensive drugs. *Int J Cardiol* 2008 Dec 5. [Epub ahead of print]
148. Piecha G, Adamczak M, Chudek J, et al. Indapamide decreases plasma adiponectin concentration in patients with essential hypertension. *Kidney Blood Press Res* 2007;30:187–94.
149. Yilmaz MI, Sonmez A, Caglar K, et al. Effect of antihypertensive agents on plasma adiponectin levels in hypertensive patients with metabolic syndrome. *Nephrology (Carlton)* 2007;12:147–53.
150. Benndorf RA, Rudolph T, Appel D, et al. Telmisartan improves insulin sensitivity in nondiabetic patients with essential hypertension. *Metabolism* 2006;55:1159–64.
151. Koh KK, Han SH, Ahn JY, et al. Amlodipine improves endothelial function and metabolic parameters in patients with hypertension. *Int J Cardiol* 2009;133: 23–31.
152. Furuhashi M, Ura N, Higashiura K, et al. Blockade of the renin–angiotensin system increases adiponectin concentrations in patients with essential hypertension. *Hypertension* 2003;42:76–81.
153. Koh KK, Quon MJ, Han SH, et al. Anti-inflammatory and metabolic effects of candesartan in hypertensive patients. *Int J Cardiol* 2006;108:96–100.
154. Delles C, Raff U, Mimran A, et al. Effects of telmisartan and ramipril on adiponectin and blood pressure in patients with type 2 diabetes. *Am J Hypertens* 2008;21: 1330–6.
155. Matsumoto S, Takebayashi K, Aso Y. The effect of spironolactone on circulating adipocytokines in patients with type 2 diabetes mellitus complicated by diabetic nephropathy. *Metabolism* 2006;55:1645–52.
156. Uchida T, Shimizu M, Sakai Y, et al. Effects of losartan on serum total and high-molecular weight adiponectin concentrations in hypertensive patients with metabolic syndrome. *Metabolism* 2008;57:1278–85.
157. Ouchi N, Kihara S, Arita Y, et al. Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation* 1999;100: 2473–6.
158. Kumada M, Kihara S, Sumitsuiji S, et al. Association of hypoadiponectinemia with coronary artery disease in men. *Arterioscler Thromb Vasc Biol* 2003;23:85–9.
159. Nakamura Y, Shimada K, Fukuda D, et al. Implications of plasma concentrations of adiponectin in patients with coronary artery disease. *Heart* 2004;90:528–33.
160. Pilz S, Maerz W, Weihrauch G, et al. Adiponectin serum concentrations in men with coronary artery disease: the Ludwigshafen Risk and Cardiovascular Health (LURIC) study. *Clin Chim Acta* 2006;364:251–5.
161. Lu G, Chiem A, Anuurad E, et al. Adiponectin levels are associated with coronary artery disease across Caucasian and African-American ethnicity. *Transl Res* 2007;149: 317–23.
162. Costacou T, Zgibor JC, Evans RW, et al. The prospective association between adiponectin and coronary artery disease among individuals with type 1 diabetes. The Pittsburgh epidemiology of diabetes complications study. *Diabetologia* 2005;48:41–8.
163. Schnabel R, Messow CM, Lubos E, et al. Association of adiponectin with adverse outcome in coronary artery disease patients: results from the atherogene study. *Eur Heart J* 2008;29:649–57.
164. Takemoto F, Katori H, Sawa N, et al. Plasma adiponectin: a predictor of coronary heart disease in hemodialysis patients—a Japanese prospective eight-year study. *Nephron Clin Pract* 2008;111:c12–20.
165. Lim S, Koo BK, Cho SW, et al. Association of adiponectin and resistin with cardiovascular events in Korean patients with type 2 diabetes: the Korean Atherosclerosis Study (KAS): a 42-month prospective study. *Atherosclerosis* 2008;196:398–404.
166. Cavadoglu E, Ruwende C, Chopra V, et al. Adiponectin is an independent predictor of all-cause mortality, cardiac mortality, and myocardial infarction in patients presenting with chest pain. *Eur Heart J* 2006;27:2300–9.
167. Pischon T, Girman CJ, Hotamisligil GS, et al. Plasma adiponectin levels and risk of myocardial infarction in men. *JAMA* 2004;291:1730–7.
168. Lawlor DA, Davey Smith G, Ebrahim S, et al. Plasma adiponectin levels are associated with insulin resistance, but do not predict future risk of coronary heart disease in women. *J Clin Endocrinol Metab* 2005;90:5677–83.
169. Soderberg S, Colquhoun D, Keech A, et al. Leptin, but not adiponectin, is a predictor of recurrent cardiovascular events in men: results from the lipid study. *Int J Obes (Lond)* 2009;33:123–30.
170. Sattar N, Wannamethee G, Sarwar N, et al. Adiponectin and coronary heart disease: a prospective study and meta-analysis. *Circulation* 2006;114:623–9.
171. Aso Y, Yamamoto R, Wakabayashi S, et al. Comparison of serum high-molecular weight (HMW) adiponectin with total adiponectin concentrations in type 2 diabetic patients with coronary artery disease using a novel enzyme-linked immunosorbent assay to detect HMW adiponectin. *Diabetes* 2006;55:1954–60.
172. Inoue T, Kotooka N, Morooka T, et al. High molecular weight adiponectin as a predictor of long-term clinical

- outcome in patients with coronary artery disease. *Am J Cardiol* 2007;100:569–74.
173. von Eynatten M, Humpert PM, Bluemm A, et al. High-molecular weight adiponectin is independently associated with the extent of coronary artery disease in men. *Atherosclerosis* 2008;199:123–8.
 174. Rizza S, Clementi F, Porzio O, et al. Adiponectin isoforms are not associated with the severity of coronary atherosclerosis but with undiagnosed diabetes in patients affected by stable CAD. *Nutr Metab Cardiovasc Dis* 2009;19:54–60.
 175. von Eynatten M, Hamann A, Twardella D, et al. Atherogenic dyslipidaemia but not total- and high-molecular weight adiponectin are associated with the prognostic outcome in patients with coronary heart disease. *Eur Heart J* 2008;29:1307–15.
 176. Sattar N, Watt P, Cherry L, et al. High molecular weight adiponectin is not associated with incident coronary heart disease in older women: a nested prospective case-control study. *J Clin Endocrinol Metab* 2008;93:1846–9.
 177. Hwang JJ, Yang WS, Chiang FT, et al. Association of circulating matrix metalloproteinase-1, but not adiponectin, with advanced coronary artery disease. *Atherosclerosis* 2008 Aug 28. [Epub ahead of print]
 178. Chen MP, Tsai JC, Chung FM, et al. Hypoadiponectinemia is associated with ischemic cerebrovascular disease. *Arterioscler Thromb Vasc Biol* 2005;25:821–6.
 179. Efstathiou SP, Tsioulos DI, Tsiakou AG, et al. Plasma adiponectin levels and five-year survival after first-ever ischemic stroke. *Stroke* 2005;36:1915–9.
 180. Soderberg S, Stegmayr B, Stenlund H, et al. Leptin, but not adiponectin, predicts stroke in males. *J Intern Med* 2004;256:128–36.
 181. Matsumoto M, Ishikawa S, Kajii E. Association of adiponectin with cerebrovascular disease: a nested case-control study. *Stroke* 2008;39:323–8.
 182. Stott DJ, Welsh P, Rumley A, et al. Adipocytokines and risk of stroke in older people: a nested case-control study. *Int J Epidemiol* 2009;38:253–61.
 183. Lim PS, Hu CY, Wu MY, et al. Plasma adiponectin is associated with ankle-brachial index in patients on haemodialysis. *Nephrology (Carlton)* 2007;12:546–52.
 184. Kokubu N, Tsuchihashi K, Yuda S, et al. Persistent insulin-sensitizing effects of sarpogrelate hydrochloride, a serotonin 2a receptor antagonist, in patients with peripheral arterial disease. *Circ J* 2006;70:1451–6.
 185. Kistorp C, Faber J, Galatius S, et al. Plasma adiponectin, body mass index, and mortality in patients with chronic heart failure. *Circulation* 2005;112:1756–62.
 186. George J, Patal S, Wexler D, et al. Circulating adiponectin concentrations in patients with congestive heart failure. *Heart* 2006;92:1420–4.
 187. Wannamethee SG, Whincup PH, Lennon L, et al. Circulating adiponectin levels and mortality in elderly men with and without cardiovascular disease and heart failure. *Arch Intern Med* 2007;167:1510–7.
 188. Tamura T, Furukawa Y, Taniguchi R, et al. Serum adiponectin level as an independent predictor of mortality in patients with congestive heart failure. *Circ J* 2007;71:623–30.
 189. Haugen E, Furukawa Y, Isic A, et al. Increased adiponectin level in parallel with increased NT-pro BNP in patients with severe heart failure in the elderly: a hospital cohort study. *Int J Cardiol* 2008;125:216–9.
 190. Norrelund H, Wiggers H, Halbirk M, et al. Abnormalities of whole body protein turnover, muscle metabolism and levels of metabolic hormones in patients with chronic heart failure. *J Intern Med* 2006;260:11–21.
 191. McEntegart MB, Awede B, Petrie MC, et al. Increase in serum adiponectin concentration in patients with heart failure and cachexia: relationship with leptin, other cytokines, and b-type natriuretic peptide. *Eur Heart J* 2007;28:829–35.
 192. Ingelsson E, Riserus U, Berne C, et al. Adiponectin and risk of congestive heart failure. *JAMA* 2006;295:1772–4.
 193. Tsutamoto T, Tanaka T, Sakai H, et al. Total and high molecular weight adiponectin, haemodynamics, and mortality in patients with chronic heart failure. *Eur Heart J* 2007;28:1723–30.
 194. Komaba H, Igaki N, Goto S, et al. Adiponectin is associated with brain natriuretic peptide and left ventricular hypertrophy in hemodialysis patients with type 2 diabetes mellitus. *Nephron Clin Pract* 2007;107:c103–8.
 195. von Eynatten M, Hamann A, Twardella D, et al. Relationship of adiponectin with markers of systemic inflammation, atherogenic dyslipidemia, and heart failure in patients with coronary heart disease. *Clin Chem* 2006;52:853–9.
 196. Tanaka T, Tsutamoto T, Sakai H, et al. Effect of atrial natriuretic peptide on adiponectin in patients with heart failure. *Eur J Heart Fail* 2008;10:360–6.
 197. Shimano M, Shibata R, Tsuji Y, et al. Circulating adiponectin levels in patients with atrial fibrillation. *Circ J* 2008;72:1120–4.